

t_k moves on links

JÓZEF H. PRZYTICKI

Abstract: It is a natural question to ask whether two links are equivalent by the following moves $\xrightarrow{t_k} \mathfrak{X} \cdots \mathfrak{X}$ (where k is a fixed number of positive half twists) and if they are, how many moves are needed to go from one link to the other. In particular if $k = 2$ and the second link is a trivial link it is the question about the unknotting number. The new polynomial invariants of links often allow us to answer the above questions. Also the first homology groups of cyclic branch covers over links provide some interesting information.

Introduction. In the first part of the paper we apply the Jones-Conway (Homfly) and Kauffman polynomials to find whether two links are not t_k equivalent and if they are, to gain some information how many moves are needed to go from one link to the other.

In the second part we describe the Fox congruence classes and their relations with t_k moves. We use the Fox method to analyse relations between t_k moves and the first homology groups of branched cyclic covers of links.

In the third part we consider the influence of t_k moves on the Goeritz and Seifert matrices and analyse Lickorish-Millett [L-M-2] and Murakami [Mur-1, Mur-2] formulas from the point of view of t_k moves and illustrate them by various examples. At the end of the paper we outline some relations with signatures of links and non-cyclic coverings of link spaces.

Now we will formulate the basic definitions and state the main results of the paper concerning connections between t_k moves and the Jones-Conway polynomial invariants of links.

Consider diagrams of oriented links L_0 and L_k which are identical, except the parts of the diagrams shown on Fig. 0.1.




Fig. 0.1

Definition 0.1. The t_k move (or k twist) is the elementary operation on an oriented diagram L_0 resulting in L_k (Fig. 0.1). Two oriented links L and L' are said to be t_k equivalent ($L \sim_{t_k} L'$) if one can go from L to L' using $t_k^{\pm 1}$ moves (and isotopy). The t_k distance between t_k equivalent links L and L' (denoted $|L, L'|_{t_k}$) is defined to be the minimal number of $t_k^{\pm 1}$ moves needed to go from L to L' .

The t_k level distance between L and L' (denoted $|L, L'|_{t_k}^{\text{lev}}$) is defined to be the number of t_k moves minus the number of t_k^{-1} moves needed when we go from L to L' (we will show later (Corollary 1.2) that for $k > 2$ it does not depend on the choice of a path joining L and L').

The classical unknotting number is the t_2 distance from a given link to an unlink.

Corollary 0.2. Let $P_L(a, z)$ be a Jones-Conway polynomial described by the properties

- (i) $P_{T_1}(a, z) = 1,$

(ii) $aP_{\nearrow\searrow}(a, z) + a^{-1}P_{\searrow\nearrow}(a, z) = zP_{\rightrightarrows}(a, z)$,
 where T_1 is a trivial knot. Then for $z_0 = 2 \cos(\pi m/k)$ ($z_0 \neq 0, \mp 2$)


$$P_{t_k(L)}(a, z_0) = (-1)^m a^{-k} P_L(a, z_0)$$

and for t_k equivalent links L and L'

$$P_{L'}(a, z_0) = ((-1)^m a^{-k})^{|L, L'|_{t_k}^{\text{lev}}} P_L(a, z_0)$$

and neither side is identically zero.

We can introduce a $\bar{t}_k$ move and $\bar{t}_k$ equivalence of oriented links ($\sim_{\bar{t}_k}$) similarly to the t_k move and ($\sim_{t_k}$) (see Fig. 0.2).



($\bar{t}_k(L)$ is naturally oriented if k is even)

Fig. 0.2

Corollary 0.3. (1.8) If $a_0^{2k} = (-1)^k$, $a_0 \neq \mp i$, then

$$P_{\bar{t}_{2k}(L)}(a_0, z) = P_L(a_0, z).$$

Corollary 0.4. Let $V_L(t)$ be the Jones polynomial described by the properties

- (i) $V_{T_1}(t) = 1$,
- (ii) $t^{-1}V_{\searrow\nearrow}(t) - tV_{\nearrow\searrow}(t) = (\sqrt{t} - \frac{1}{\sqrt{t}})V_{\rightrightarrows}(t)$,

then

- (a) If $t^k = (-1)^k$ (i.e. $t^{1/2} = -ie^{\pi im/k}$), $t \neq -1$, then a t_k move changes $V_L(t)$ by $(-1)^m i^k$, that is

$$V_{t_k(L)}(t) = (-1)^m i^k V_L(t).$$

- (b) If $t^{2k} = 1$ (i.e. $t = e^{\pi im/k}$), $t \neq -1$, then

$$V_{\bar{t}_{2k}(L)}(t) = V_L(t).$$

- (c) Assume k is odd and $t^k = -1$. Then

$$V_{\bar{t}_k(L)}(t) = \omega_{4k} V_L(t),$$

where ω_{4k} is a properly chosen $4k$ -root of unity (depending also on the choice of the orientation of $\bar{t}_k(L)$; see Theorem 1.13).

1. t_k -MOVES AND CONWAY FORMULAS FOR THE JONES-CONWAY AND KAUFFMAN POLYNOMIALS.

When one considers the sequence of links $L, t_1(L), t_2(L), \dots, (\asymp, \asymp, \asymp, \dots)$ then the Jones-Conway (and Kauffman) polynomials $P_L(a, z), P_{t_1(L)}(a, z), P_{t_2(L)}(a, z), \dots$ form a (generalized) Fibonacci sequence. So one can expect that there is a nice formula which expresses $P_{t_k(L)}(a, z)$ in terms of $P_{t_1(L)}(a, z)$ and $P_L(a, z)$ and in fact we have the following result:

Theorem 1.1. $a^k P_{t_k(L)}(a, z) = a v_1^{(k)}(z) P_{t_1(L)}(a, z) - v_1^{(k-1)}(z) P_L(a, z)$, where $v_1^{(k+2)}(z) = z v_1^{(k+1)}(z) - v_1^{(k)}(z)$ and $v_1^{(-1)}(z) = -1$, $v_1^{(0)}(z) = 0$, $v_1^{(1)}(z) = 1$. In particular if one substitutes $z = p + p^{-1}$ one gets $v_1^{(k)}() = \frac{p^k - p^{-k}}{p - p^{-1}}$. [Added for e-print: $v_1^{(k)}(z)$ is a variant of the Chebyshev polynomial of the second kind.]

Proof. We proceed by induction on k . For $k = 1, 2$ the formula from Theorem 1.1 holds:

$$a P_{t_1(L)}(a, z) = a P_{t_1(L)}(a, z) - 0 \cdot P_L(a, z)$$

and

$$a^2 P_{t_2(L)}(a, z) = az P_{t_1(L)}(a, z) - P_L(a, z).$$

Assume that Theorem 1.1 holds for $1, 2, \dots, k-1$, ($k > 2$). Now one gets:

$$\begin{aligned} a^k P_{t_k(L)}(a, z) &= a^{k-1} z P_{t_{k-1}(L)}(a, z) - a^{k-2} z P_{t_{k-2}(L)}(a, z) = \\ &= z(av_1^{(k-1)}(z)P_{t_1(L)}(a, z) - v_1^{(k-2)}(z)P_L(a, z)) - \\ &\quad (av_1^{(k-2)}(z)P_{t_1(L)}(a, z) - v_1^{(k-3)}(z)P_L(a, z)) = \\ &= a(zv_1^{(k-1)}(z)P_{t_1(L)}(a, z) - v_1^{(k-2)}(z)P_{t_1(L)}(a, z)) - \\ &\quad (zv_1^{(k-2)}(z)P_L(a, z) - v_1^{(k-3)}(z)P_L(a, z)) = \\ &= av_1^{(k)}(z)P_{t_1(L)}(a, z) - v_1^{(k-1)}(z)P_L(a, z). \end{aligned}$$

To see, that for $z = p + p^{-1}$, $v_1^{(k)}(z) = \frac{p^k - p^{-k}}{p - p^{-1}}$ it is enough to observe that

$$\frac{p^{k+2} - p^{-(k+2)}}{p - p^{-1}} = (p + p^{-1}) \frac{p^{k+1} - p^{-(k+1)}}{p - p^{-1}} - \frac{p^k - p^{-k}}{p - p^{-1}}.$$

□

Corollary 1.2. *If $p_0^{2k} = 1$ (i.e. $p_0 = e^{\pi i m/k}$), $p_0 \neq \pm 1, \pm i$ or equivalently $z_0 = 2 \cos(\pi m/k)$; $z_0 \neq 0, \pm 2$ then*

$$P_{t_k(L)}(a, z_0) = (-1)^m a^{-k} P_L(a, z_0)$$

and for t_k equivalent links L and L'

$P_{L'}(a, z_0) = ((-1)^m a^{-k})^{|L, L'|_{t_k}^{\text{lev}}} P_L(a, z_0)$ and neither side is identically zero.

Proof. Assume $p_0 \neq \mp 1, \mp i$. Then $v_1^{(k)}(z_0) = 0$ reduces to $p_0^{2k} = 1$, so $p_0 = e^{\pi im/k}$ and $z_0 = p_0 + p_0^{-1} = 2 \cos(\pi m/k)$. Now the equation from Theorem 1.1 reduces to

$$\begin{aligned} a^k P_{t_k(L)}(a, z_0) &= -v_1^{(k-1)}(z_0) P_L(a, z_0) = -\frac{p_0^{k-1} - p_0^{1-k}}{p_0 - p_0^{-1}} P_L(a, z_0) = \\ &= p_0^k P_L(a, z_0) = (-1)^m P_L(a, z_0). \end{aligned}$$

So the first part of Corollary 1.2 is proven.

For the second part it is enough to show that for each link L and any complex number z_0 ($z_0 \neq 0$) $P_L(a, z_0)$ is never identically zero. It follows from the fact that $P_L(a, a + a^{-1}) \equiv 1$ (see [L-M-1] or [P-1], or apply the standard induction: it holds for trivial links and whenever it holds for $L_-(\nearrow)$ and $L_0(\rightrightarrows)$ it holds for $L_+(\searrow)$ and if it holds for L_+ and L_0 it holds for L_-). $\square$

If $a = i$, $t^{1/2} = -ip$ in the Jones-Conway polynomial $P_L(a, z)$, ($z = p + p^{-1}$), we get the (normalized) Alexander polynomial $\Delta_L(t)$ which satisfies:

- (i) $\Delta_{T_1}(t) = 1$,
- (ii) $\Delta_{L_+}(t) - \Delta_{L_-}(t) = (\sqrt{t} - \frac{1}{\sqrt{t}}) \Delta_{L_0}(t)$.

Corollary 1.3. [Fo-1, Ki] *If $t^k = (-1)^k$ (i.e. $t^{1/2} = -ie^{\pi im/k}$), $t \neq -1$ then $\Delta_{t_k(L)}(t) = (-1)^m (-i)^k \Delta_L(t)$.*

Proof. It follows immediately from Corollary 1.2. One have only additionally notice that the formula from Corollary 1.2 remains true for $a = \mp i$, $p = \mp i$. $\square$

When we substitute $a = it^{-1}$, $p = it^{1/2}$ in $P(a, z)$ ($z = p + p^{-1}$) we get the Jones polynomial $V_L(t)$ which satisfies:

- (i) $V_{T_1}(t) = 1$,
- (ii) $\frac{1}{t} V_{L_+}(t) - t V_{L_-}(t) = (\sqrt{t} - \frac{1}{\sqrt{t}}) V_{L_0}(t)$.

There has been some confusion as to the conventions. We use that of [Jo-2].

Corollary 1.4. *If $t^k = (-1)^k$ (i.e. $t^{1/2} = -ie^{\pi im/k}$), $t \neq -1$, then a t_k move changes $V_L(t)$ by $(-1)^m i^k$ that is*

$$V_{t_k(L)}(t) = (-1)^m i^k V_L(t).$$

Proof. It is true for $t = 1$ (then k is even). For $t \neq \mp 1$ it follows immediately from Corollary 1.2. $\square$

Corollary 1.5. *If $p_0 = \varepsilon = \mp 1$ (so $z_0 = 2\varepsilon = \mp 2$) then*

- (a) $a^k P_{t_k(L)}(a, z_0) = \varepsilon k a P_{t_1(L)}(a, z_0) - \varepsilon(k-1) P_L(a, z_0)$ and
- (b) $P_{t_k(L)}(a, z_0) \equiv \varepsilon a^{-k} P_L(a, z_0) \pmod{k/2^i}$ i.e. the equality holds if $P_L(a, z_0)$ is understood to be a Laurent polynomial in a with coefficients in the ring $\mathbf{Z}[1/2]/k\mathbf{Z}[1/2]$.

Corollary 1.6. *(Generalized Conway formula). The following formula holds for the Jones-Conway polynomial:*

$$a^k P_{L_{+k}}(a, z) + a^{-k} P_{L_{-k}}(a, z) = w_1^{(k)}(z) P_{L_0}(a, z),$$

where $w_1^{(0)} = 2$, $w_1^{(1)} = z$, $w_1^{(k)} = z w_1^{(k-1)} - w_1^{(k-2)}$. After substituting $z = p + p^{-1}$ one gets $w_1^{(k)} = p^k + p^{-k}$.

Proof. From Theorem 1.1 one gets:

$$a^k P_{L_{+k}} = a v_1^{(k)} P_{L_{+1}} - v_1^{(k-1)} P_{L_0}$$

and

$$a^{-k} P_{L_{-k}} = a^{-1} v_1^{(k)} P_{L_{-1}} - v_1^{(k-1)} P_{L_0}.$$

Adding these equations by sides one gets:

$$a^k P_{L_{+k}} + a^{-k} P_{L_{-k}} = v_1^{(k)} (a P_{L_{+1}} + a^{-1} P_{L_{-1}}) - 2 v_1^{(k-1)} P_{L_0} =$$

$$= (zv_1^{(k)} - 2v_1^{(k-1)})P_{L_0}.$$

Now substituting $w_1^{(k)} = zv_1^{(k)} - 2v_1^{(k-1)}$ one gets the equation from Corollary 1.6 (notice that $v_1^{(-1)} = -1$). $\square$

Now we will get formulas for $\bar{t}_k$ moves analogous to those for t_k moves.

Theorem 1.7. $P_{\bar{t}_{2k}(L)}(a, z) = (-1)^k a^{2k} P_{L \asymp}(a, z) + zu_1^{(2k)}(a) P_{L \wr}(a, z)$ where $\bar{t}_{2k}(L)$, $L \asymp$ and $L \wr$ are oriented diagrams which are identical, except the parts of the diagrams shown on Fig. 1.1, and $u_1^{(0)} = 0$, $u_1^{(2)} = a$, $u_1^{(2k)} = -a^2 u_1^{(2(k-1))} + a$ or equivalently $u_1^{(2k)} = (-1)^{k+1} a^k \frac{a^k + (-1)^{k+1} a^{-k}}{a + a^{-1}}$.




Fig. 1.1

Proof. We proceed by induction on k . For $k = 0, 1$ the formula from Theorem 1.7 holds:

$$P_{L \asymp}(a, z) = P_{L \asymp}(a, z) + z \cdot 0 \cdot P_{L \wr}(a, z)$$

and

$$P_{\bar{t}_2(L)}(a, z) = -a^2 P_{L \asymp}(a, z) + za P_{L \wr}(a, z).$$

Assume that Theorem 1.7 holds for $0, 1, \dots, k-1$ ($k \geq 2$). Now one gets:

$$\begin{aligned}
 P_{\bar{t}_{2k}(L)} &= -a^2 P_{t_{2(k-1)}(L)} + za P_L \text{ (crossing)} = -a^2 ((-1)^{k-1} a^{2(k-1)} P_L \text{ (cup)}) \\
 &\quad -a^2 (zu_1^{(2(k-1))} P_L \text{ (crossing)} + za P_L \text{ (crossing)} = (-1)^k a^{2k} P_L \text{ (cup)} + \\
 &\quad z(-a^2 u_1^{(2(k-1))} + a) P_L \text{ (crossing)} = (-1)^k a^{2k} P_L \text{ (cup)} + zu_1^{(2k)} P_L \text{ (crossing)}.
 \end{aligned}$$

□

Corollary 1.8. *If $a_0^{2k} = (-1)^k$ ($a_0 \neq \mp i$), then*

$$P_{\bar{t}_{2k}(L)}(a_0, z) = P_L(a_0, z).$$

Corollary 1.9. *If $a_0 = \varepsilon i = \mp i$ then*

- (a) $P_{\bar{t}_{2k}(L)}(a_0, z) = P_L \text{ (cup)}(a_0, z) + z\varepsilon i k P_L \text{ (crossing)}(a_0, z)$ and
- (b) $P_{\bar{t}_{2k}(L)}(a_0, z) \equiv P_L(a_0, z) \pmod{k}$ i.e. equality holds if $P_L(a_0, z)$ is understood to be the Laurent polynomial in z with coefficients in the ring $\mathbf{Z} + i\mathbf{Z}/k(\mathbf{Z} + i\mathbf{Z})$.
- (c) [Fo-1] If $a = i$, $t^{1/2} = -ip$ ($z = p + p^{-1}$) one gets the (normalized) Alexander polynomial and (b) reduces to $\Delta_{\bar{t}_{2k}(L)}(t) \equiv \Delta_L(t) \pmod{k}$ i.e. the equality holds if $\Delta_L(t)$ is reduced to a polynomial in $\mathbf{Z}_k[\sqrt{t}^{\mp 1}]$.

For the Jones polynomial ($a = it^{-1}$, $p = it^{1/2}$), Corollary 1.8 reduces to:

Corollary 1.10. *If $t^{2k} = 1$, $t \neq -1$ then*

$$V_{\bar{t}_{2k}(L)}(t) = V_L(t).$$

Proof. It is true for $t = 1$. For $t \neq \mp 1$ it follows from Corollary 1.8. □

Corollary 1.11. (*Generalized Conway formula*). *The following formulas hold for the Jones-Conway polynomial:*

- (i) $a^{-2k}P_{\bar{t}_{2k}(L)}(a, z) + a^{2k}P_{\bar{t}_{2k}^{-1}(L)}(a, z) = (-1)^k 2P_{L_{\bigcap}}(a, z) + z\left(\frac{a^k + (-1)^{k+1}a^{-k}}{a + a^{-1}}\right)^2 P_{L_{\bigcap}}(a, z),$
- (ii) $a^{-k}P_{t_{2k}(L)}(a, z) + (-1)^k a^k P_{\bar{t}_{2k}^{-1}(L)}(a, z) = ((-1)^k a^k + a^{-k})P_{L_{\bigcap}}(a, z),$
- (iii) $a^{-k}P_{t_k(L)}(a, z) + (-1)^{k+1}a^k P_{\bar{t}_k^{-1}(L)}(a, z) = z\left(\frac{a^{-k} + \varepsilon(k)a^k}{a + a^{-1}}\right)P_{L_{\bigcap}}(a, z),$

where

$$\varepsilon(k) = \begin{cases} -1 & \text{if } k+2 \text{ is a multiple of } 4, \\ 1 & \text{otherwise} \end{cases}$$

Proof. (i) follows immediately from Theorem 1.7; one has to add equations for $a^{-2k}P_{\bar{t}_{2k}(L)}$ and for $a^{2k}P_{\bar{t}_{2k}^{-1}(L)}$.

(ii) follows from Theorem 1.7, by adding equations for $a^{-k}P_{t_{2k}(L)}$ and for $(-1)^k a^k P_{\bar{t}_{2k}^{-1}(L)}$.

(iii) (k even) follows from Theorem 1.7 by adding equations for $a^{-2k}P_{t_k(L)}$ and for $(-1)a^{2k}P_{\bar{t}_k^{-1}(L)}$. If k is odd then from Theorem 1.7 one gets

$$a^{-(2k+1)}P_{\bar{t}_{2k+1}(L)}(a, z) = (-1)^k a^{-1}P_{\bar{t}_1(L)} + za^{-(2k+1)}u_1^{(2k)}(a)P_{L_{\bigcap}}(a, z)$$

and

$$a^{2k+1}P_{\bar{t}_{2k+1}^{-1}(L)}(a, z) = (-1)^k aP_{\bar{t}_1^{-1}(L)}(a, z) + za^{2k+1}((-1)^{k+1}a^{-2k}u_1^{(2k)}(a))P_{L_{\bigcap}}(a, z)$$

(in the last equality we use the fact that $u_1^{(2k)}(1/a) = (-1)^{k+1}a^{-2k}u_1^{(2k)}(a)$)

Adding the above equalities one gets:

$$\begin{aligned} & a^{-2k+1}P_{\bar{t}_{2k+1}(L)} + a^{2k+1}P_{\bar{t}_{2k+1}^{-1}(L)} = (-1)^k(a^{-1}P_{\bar{t}_1(L)} + aP_{\bar{t}_1^{-1}(L)}) + \\ & + (z(-1)^{k-1}a^{-(k+1)}\frac{a^k + (-1)^{k+1}a^{-k}}{a + a^{-1}} + za^{k+1}\frac{a^k + (-1)^{k+1}a^{-k}}{a + a^{-1}})P_{L_{\bigcap}} = \end{aligned}$$

$$\begin{aligned} zP_L \cdot ((-1)^k + \frac{(-1)^{k+1}a^{-1} + a^{-(2k+1)} + a^{2k+1} + (-1)^{k+1}a}{a + a^{-1}}) = \\ = z(\frac{a^{2k+1} + a^{-(2k+1)}}{a + a^{-1}})P_L. \end{aligned}$$

□

We worked, till now, with $\bar{t}_k$ moves for k even, and the reason for this was that if L is oriented then $\bar{t}_k(L)$ has no any natural orientation for k odd. For the Jones polynomial, however, one has Jones reversing result (see [L-M-2] or [P-1]) so one can still find how $V_L(t)$ is changed under a $\bar{t}_k$ move.

Namely let $L = \{L_1, \dots, L_i, \dots, L_n\}$ be an oriented link of n components and $L' = \{L_1, \dots, -L_i, \dots, L_n\}$, i.e. the orientation of L_i is reversed, and let $\lambda = lk(L_i, L - L_i)$. Then

$$\mathbf{1.12.} \quad V_{L'}(t) = t^{-3\lambda} V_L(t).$$

Theorem 1.13. *Consider a $\bar{t}_k$ move on an oriented link L , and assume k is odd. We have two cases:*

- (i) $c(\bar{t}_k(L)) < c(L)$, where $c(L)$ denotes the number of components. That is two components of L , say L_i and L_j , are involved in the $\bar{t}_k$ move (see Fig.1.2). Let $\lambda = lk(L_i, L - L_i)$. Then for $t^k = (-1)^k$ (i.e. $t^{1/2} = -ie^{\pi im/k}$), $i \neq -1$:

$$V_{\bar{t}_k(L)}(t) = (-1)^m i^k t^{-3\lambda} V_L(t) = (-1)^{m+\lambda} i^k e^{-6\pi im\lambda/k} V_L(t),$$

where the orientation of $\bar{t}_k(L)$ is chosen so that it does not agree with the orientation of L_i .

- (ii) $c(\bar{t}_k(L)) = c(L)$. That is one component of L is involved in the $\bar{t}_k$ move. Let $L_{\smile}$ denote the smoothing of L (Fig. 1.3). $L_{\smile}$ has more components than L and let L_i, L_j be the new components of $L_{\smile}$ (Fig. 1.3). Let $\lambda = lk(L_i, L_{\smile} - L_i)$ and assume that $\bar{t}_k(L)$ is oriented in such a way that its orientation agrees with that of $L_{\smile}$ with exception of L_i . Then for $t^k = (-1)^k$ (i.e. $t^{1/2} = -ie^{\pi im/k}$), $i \neq -1$:

$$V_{\bar{t}_k(L)}(t) = t^{-3\lambda} V_L(t) = (-1)^\lambda e^{-6\pi i m \lambda / k} V_L(t).$$

Proof. (i) We use the Jones reversing result and Corollary 1.4 and we get (see Fig. 1.2)

$$V_{\bar{t}_k(L)}(t) = (-1)^m i^k V_{L'}(t) = (-1)^m i^k t^{-3\lambda} V_L(t) = (-1)^{m+\lambda} i^k e^{6\pi i m \lambda / k} V_L(t).$$




Fig. 1.2

(ii) We use Corollary 1.4 and the part (i) and we get (see Fig. 1.4):

$$V_{\bar{t}_k(L)}(t) = (-1)^m i^k t^{-3\lambda+k} V_{L'}(t) = (-1)^m i^k t^{-3\lambda+k} (-1)^m i^k V_L(t) = t^{-3\lambda} V_L(t) =$$

$$(-1)^\lambda e^{-6\pi i m \lambda / k} V_L(t).$$




Fig. 1.3




Fig. 1.4

□

It is possible to get Theorem 1.13 by considering the variant of the Jones polynomial which is an invariant of regular isotopy and does not depend on orientation ([Ka-4]).

We will use this idea considering how the Kauffman polynomial changes under t_k and $\bar{t}_k$ moves.

Two diagrams of links are regularly isotopic iff one can be obtained from the other by a sequence of Reidemeister moves of type $\Omega_2^{\mp 1}$, $\Omega_3^{\mp 1}$ and isotopy of the projection plane (see Fig.1.5).




Fig. 1.5

The Kauffman polynomial of regular isotopy of unoriented diagrams is defined by (see [Ka-3]; also [P-1]):

- (1) $\Lambda_{T_1}(a, x) = a^{\text{tw}(T_1)}$, where T_1 is a diagram representing the trivial knot (up to isotopy) and $\text{tw}(T_1) = \sum \text{sgn } p$ where the sum is taken over all crossings of T_1 .
- (2) $\Lambda_{\nearrow \searrow}(a, x) + \Lambda_{\searrow \nearrow}(a, x) = x\Lambda_{\text{=}}(a, x) + x\Lambda_{\text{)} \text{ (}}(a, x)$.

The Kauffman polynomial of oriented links is defined by

$$F_L(a, x) = a^{-\text{tw}(L)} \Lambda_L(a, x).$$

Theorem 1.14. $\underbrace{\Lambda_{\infty \dots \infty}}_{k \text{ half twists}}(a, x) = v_1^{(k)}(x) \Lambda_{\nearrow \searrow}(a, x) - v_1^{(k-1)}(x) \Lambda_{\text{=}}(a, x) + x v_2^{(k)}(a, x) \Lambda_{\text{)} \text{ (}}(a, x)$, k half twists where $v_1^{(k)}()$ is the same as in Theorem 1.1 and $v_2^{(0)}(a, x) = 0$, $v_2^{(1)}(a, x) = 0$, $v_2^{(2)}(a, x) = a^{-1}$, $v_2^{(k)}(a, x) = x v_2^{(k-1)}(a, x) - v_2^{(k-2)}(a, x) + a^{1-k}$. In particular for $x = p + p^{-1}$ one gets $v_1^{(k)}() = \frac{p^k - p^{-k}}{p - p^{-1}}$, $v_2^{(k)} = ((p - p^{-1})(a + a^{-1} - (p + p^{-1})))^{-1}(-a^{-1}(p^k - p^{-k}) + p(a^{-k} - p^{-k}) - p^{-1}(a^{-k} - p^k))$.

Proof. We proceed by induction on k . For $k = 0, 1, 2$ the formula from Theorem 1.14 holds. Assume that it holds for $0, 1, \dots, k-1$ ($k > 2$). Now one gets:

$$\begin{aligned}
 \underbrace{\Lambda_{\infty \dots \infty}}_{k \text{ half twists}} &= \underbrace{x \Lambda_{\infty \dots \infty}}_{(k-1) \text{ half twists}} - \underbrace{\Lambda_{\infty \dots \infty}}_{(k-2) \text{ half twists}} + xa^{1-k} \Lambda_{\text{J}} (= \\
 &= x(v_1^{(k-1)} \Lambda_{\text{X}} - v_1^{(k-1)} \Lambda_{\text{=}} + xv_2^{(k-1)} \Lambda_{\text{J}}) - (v_1^{(k-2)} \Lambda_{\text{X}} - v_1^{(k-3)} \Lambda_{\text{=}} + xv_2^{(k-2)} \Lambda_{\text{J}}) + xa^{1-k} \Lambda_{\text{J}} = \\
 &= (xv_1^{(k-1)} - v_1^{(k-2)}) \Lambda_{\text{X}} - (xv_1^{(k-2)} - v_1^{(k-3)}) \Lambda_{\text{=}} + x(xv_2^{(k-1)} - v_2^{(k-2)} + a^{1-k}) \Lambda_{\text{J}} = \\
 &= v_1^{(k)} \Lambda_{\text{X}} - v_1^{(k-1)} \Lambda_{\text{=}} + xv_2^{(k)} \Lambda_{\text{J}}.
 \end{aligned}$$

The formula for $v_2^{(k)}(a, p + p^{-1})$ may be verified directly but we omit this tedious task by considering the trivial links of Fig. 1.6. From this figure we get immediately that

$$a^{-k} = \frac{p^k - p^{-k}}{p - p^{-1}} a^{-1} - \frac{p^{k-1} - p^{-(k-1)}}{p - p^{-1}} + (p + p^{-1}) v_2^{(k)}(a, x) \frac{a + a^{-1} - (p + p^{-1})}{p + p^{-1}},$$

and it finishes the proof of Theorem 1.14. $\square$



Fig. 1.6

Corollary 1.15. (a) If $p_0^{2k} = 1$ (i.e. $p_0 = e^{\pi i m/k}$), $p_0 \neq \mp 1$, $\mp i$ or equivalently $x_0 = 2 \cos(\pi m/k)$, $x_0 \neq 0, \mp 2$, then

$$\Lambda_{\infty \dots \infty} (a, x_0) = (-1)^m \Lambda_{\infty \dots \infty} (a, x_0) + \frac{a^{-k} - p_0^{-k}}{(a - p_0)(1 - a^{-1}p_0^{-1})} \Lambda_{\infty \dots \infty} (a, x_0).$$

(b) If $p_0^{2k} = 1$, $p_0 \neq \mp 1$, $\mp i$, $a_0^k = p_0^k$, $a_0 \neq p_0^{\mp 1}$ then

$$\Lambda_{\infty \dots \infty} (a_0, x_0) = (-1)^m \Lambda_{\infty \dots \infty} (a_0, x_0).$$

(c) If $p_0 = \varepsilon = \mp 1$ (so $x_0 = 2\varepsilon = \mp 2$) and $a_0^k = \varepsilon^k$, $a_0 \neq \varepsilon$ then

$$\Lambda_{\infty \dots \infty} (a_0, x_0) \equiv \varepsilon \Lambda_{\infty \dots \infty} (a_0, x_0) \pmod{k/2^i},$$

i.e. equality holds in the ring $\mathbf{Z}[a_0/2]/k\mathbf{Z}[a_0/2]$.

Proof. It follows from Theorem 1.14 similarly as Corollaries 1.2 and 1.5 followed from Theorem 1.1. $\square$

Corollary 1.16. (Generalized Conway formula)

$$\underbrace{\Lambda_{\infty \dots \infty} (a, x) + \Lambda_{\infty \dots \infty} (a, x)}_{k \text{ half twists}} = w_1^{(k)} \Lambda_{\infty \dots \infty} (a, x) + x w_2^{(k)} (a, x) \Lambda_{\infty \dots \infty} (a, x),$$

where $w_1^{(k)}(x) = w_1^k(z)$ from Corollary 1.6, and

$$w_2^{(0)}(a, x) = 0, \quad w_2^{(1)}(a, x) = 1, \quad w_2^{(k)}(a, x) = x w_2^{(k-1)}(a, x) - w_2^{(k-2)}(a, x) + a^{k-1} + a^{1-k};$$

when one substitutes $x = p + p^{-1}$ then

$$w_1^{(k)}(x) = p^k + p^{-k} \text{ and}$$

$$w_2^{(k)}(a, x) = a^{-(k-1)} p^{-(k-1)} \left(\frac{a^k - p^k}{a - p} \right) \left(\frac{1 - a^k p^k}{1 - ap} \right).$$

Proof. From Theorem 1.14 one gets:

$$\Lambda_{\infty \dots \infty} = v_1^{(k)} \Lambda_{\nearrow} - v_1^{(k-1)} \Lambda_{\overline{\overline{}}} + xv_2^{(k)}(a, x) \Lambda_{\lrcorner}$$

and

$$\Lambda_{\searrow \dots \searrow} = v_1^{(k)} \Lambda_{\searrow} - v_1^{(k-1)} \Lambda_{\overline{\overline{}}} + xv_2^{(k)}(a^{-1}, x) \Lambda_{\lrcorner}$$

Adding the above equations by sides one gets:

$$\begin{aligned} \Lambda_{\infty \dots \infty} + \Lambda_{\searrow \dots \searrow} &= v_1^{(k)} (\Lambda_{\nearrow} + \Lambda_{\searrow}) - 2v_1^{(k-1)} \Lambda_{\overline{\overline{}}} + \\ &+ x(v_2^{(k)}(a, x) + v_2^{(k)}(a^{-1}, x)) \Lambda_{\lrcorner} = (xv_1^k - 2v_1^{(k-1)}) \Lambda_{\overline{\overline{}}} + \\ &+ x(v_1^{(k)} + v_2^{(k)}(a, x) + v_2^{(k)}(a^{-1}, x)) \Lambda_{\lrcorner}. \end{aligned}$$

Now substituting $w_1^{(k)} = xv_1^k - 2v_1^{(k-1)}$ and $w_2^{(k)} = v_1^{(k)}(a) + v_2^{(k)}(a, x) + v_2^{(k)}(a^{-1}, x)$ one gets the equation from Corollary 1.16. $\square$

We end this part of the paper by translating Corollary 1.15(b) into the Kauffman polynomial of oriented links.

Corollary 1.17. *If $p_0^{2k} = 1$, $p_0 \neq \mp 1$, $\mp i$ and $a_0^k = p_0^k$, $a_0 \neq p_0^{\mp 1}$ then*

$$(a) \quad F_{\infty \dots \infty}(a_0, p_0) = a_0^k a_0^{\text{tw}(\overline{\overline{}}) - \text{tw}(\infty \dots \infty)} F_{\overline{\overline{}}}(a_0, p_0);$$

In particular

$$(b) \quad F_{t_k(L)}(a_0, p_0) = F_L(a_0, p_0),$$

$$(c) \quad F_{\bar{t}_k(L)}(a_0, p_0) = F_L(a_0, p_0),$$

$$(d) \quad F_{\bar{t}_k(L)}(a_0, p_0) = a^{4\lambda} F_L(a_0, p_0), \text{ where } k \text{ is odd and } \lambda \text{ defined as follows (compare Theorem 1.13):}$$

- (i) *If L has more components than $\bar{t}_k(L)$ and L_i is the only component of L such that the chosen orientation on $\bar{t}_k(L)$ does not agree with that of L_i then $\lambda = lk(L_i, L - L_i)$ (compare Fig. 1.2).*

- (ii) If L has the same number of components as $\bar{t}_k(L)$, consider the smoothing $L_{\mathbf{j}}$ of L (Fig. 1.3). Let L_i be the only component of L such that the chosen orientation on $\bar{t}_k(L)$ does not agree with that of L_i then $\lambda = lk(L_i, L - L_i)$ (compare Fig. 1.3).

Proof. (a) follows immediately from Corollary 1.15(b) and the definition of $F_L(a, x)$.
 (b) and (c) hold because in these cases $\text{tw}(\text{---}) - \text{tw}(\text{---}) = \mp k$;

$$(d) \text{ tw}(\text{---}) - \text{tw}(\text{---}) = \begin{cases} 4\lambda - k & \text{in the case (i)} \\ 4\lambda + k & \text{in the case (ii)} \end{cases}$$

so the equality (d) holds.

□

When one substitutes $a = t^{-3/4}$, $x = -(t^{1/4} + t^{-1/4})$ in the Kauffman polynomial $F(a, x)$ one gets the Jones polynomial $V(t)$ ([Li], see also [P-1]). Corollary 1.15 gives, therefore, some information about the behaviour of $V(t)$ under t_k and $\bar{t}_k$ moves. It happens, however, that one gets no new information comparing with Corollaries 1.4, 1.10, and Theorem 1.13.

Theorem 1.1 and 1.14 can be stated as one theorem if one uses the three variable polynomial $J_L(a, x, z)$ which generalizes the Jones-Conway and Kauffman polynomials (see [P-1]), however, one cannot gain any new information from this approach.

2. HISTORICAL BACKGROUND (FOX CONGRUENCE CLASSES).

The unknotting number of a knot was considered probably before knot theory became a science. It was a natural question to ask how many times one has to "cheat" to get from a knot an unknot. K. Reidemeister wrote in 1932 in his book [Re]: "It is very easy to define a number of knot invariants so long as one is not concerned with giving algorithms for their computation ... One can change each knot projection into projection of circle by reversing the overcrossings and undercrossings at, say, k double points of the projection. The minimal number $u(K)$ of these operations, that is, the minimal number of self-piercings, by which a knot is transformed into a circle, is a natural measure of knottedness".

The first interesting results about unknotting number were found by H. Wendt [We] in 1937. Namely Wendt proved that if $u(K)$ is the unknotting number of K and e_s is the minimal number of generators of the group $H_1(M_K^{(s)}, \mathbb{Z})$, where $M_K^{(s)}$ is the cyclic, s -fold branched cover of (S^3, K) then

$$e_s \leq u(K)(s - 1).$$

t_k and $\bar{t}_k$ moves ($\text{= } \rightarrow \text{ } \infty \dots \infty$) appear to have been first explicitly considered by S. Kinoshita in 1957 [Kin-1]. who observed that the Wendt inequality is also valid if we allow all t_{2k} and $\bar{t}_{2k}$ moves, not only t_2 moves (see Corollary 2.6(b)). The following year, 1958, R. Fox [Fo-1] considered twists of knots and congruence of knots modulo (n, q) ; the notion which is closely related, and in some sense more general, than t_{2k} and $\bar{t}_{2k}$ moves. Congruence modulo (n, q) was chosen so that the Alexander polynomial (or more generally Alexander module) is a good tool to study this.

The same year (1958), S. Kinoshita [Kin-2] used the Fox twists to generalize once more the Wendt inequality (see Corollary 2.6). The Fox approach is related to ours so we will present it here with some details. We follow the Fox paper [Fo-1] taking into account the corrections made by Kinoshita [Kin-3] and Nakanishi and Suzuki [N-S]. I am grateful to K. Murasugi and H. Murakami for informing me about the Fox paper and about the Kawauchi and Nakanishi conjectures.

Consider the following homeomorphism τ of a 3-disk $D^3 = \langle 0, 1 \rangle \times D^2$: $\tau(t, z) = (t, e^{2\pi i t} z)$. It is the natural extension to $\langle 0, 1 \rangle \times D^2$ of the Dehn twist on the annulus $\langle 0, 1 \rangle \times \partial D^2$ (see Fig. 2.1).




Fig. 2.1

We call τ a simple twist or a Dehn twist. Now whenever we have a properly embedded 2-disk in a 3-manifold M (and either M or a tubular neighbourhood of the disk is oriented), we have uniquely (up to isotopy) associated with the disk the Dehn twist (the twist is carried by a tubular neighbourhood of the disk). In particular for an oriented solid torus there is only one nontrivial Dehn twist, because there is only one, up to isotopy, nontrivial proper disk in it.

Now let L be a link in Σ^3 (we will assume $\Sigma^3 = S^3$, but in fact Σ^3 can be any homology 3-sphere), and D^2 a disk which cuts L transversely. Let V_2 be the solid torus - a small tubular neighbourhood of ∂D^2 in Σ^3 , and V_1 the closure of its complement ($V_1 = \overline{\Sigma^3 - V_2}$). If $\Sigma^3 = S^3$, V_1 is a solid torus too. Now perform the Dehn twist on V_1 using the disk D^2 . The twist restricted to the link L is denoted by $t_{2,q}$ where $q \geq 0$ is the absolute value of the crossing number of D^2 and L . By $t_{2n,q}$ we denote $t_{2,q}^n$. Notice that our t_{2n} move is special case of $t_{2n,2}$ move, and $\bar{t}_{2n}$ move is a special case of $t_{2n,0}$ move. Two oriented links L_1 and L_2 are called, by Fox, *congruent modulo n, q* ($L_1 \equiv L_2 \pmod{n, q}$) if one can go from L_1 to L_2 using $t_{2n,q'}^{\mp 1}$ moves (and isotopy), where q' can vary but is always a multiple of q . If we allow only

$t_{2n,q}^{\mp 1}$ moves then we say, after Nakanishi and Suzuki, that L_1 and L_2 are q -congruent modulo n ($L_1 \equiv_q L_2 \pmod{n}$) or that they are $t_{2n,q}$ equivalent ($L_1 \sim_{t_{2n,q}} L_2$).


The Alexander polynomial (and module) is a nice tool for distinguishing nonequivalent links because L and $t_{2n,q}(L)$ are the same outside the ball in which the move occurs.

Theorem 2.1. (a) $t_{2n,q}$ equivalent links have the same Alexander module modulo $\frac{(t-1)(t^{nq}-1)}{t^q-1}$, in particular
 (b) for $t_0^{nq} = 1$ ($t_0^q \neq 1$ or $t_0 = 1$) $\Delta_L(t) \equiv \Delta_{t_{2n,q}(L)}(t)$.

It can be understood as follows: $\Delta_L(t)$ and $\Delta_{t_{2n,q}(L)}(t)$ are equal as elements of the ring $R = \mathbf{Z}[t^{\pm 1}]/(t-1)(1+t^q+\dots+t^{(n-1)q})$, up to multiplication by invertible elements of R (in fact up to multiplication by classes of invertible elements in $\mathbf{Z}[t^{\pm 1}]$ i.e. $\mp t^p$).

If we substitute $a = i$ and $p = it^{1/2}$ in the Jones-Conway polynomial then we get the (normalized) Alexander polynomial $\Delta(t) \in \mathbf{Z}[t^{\pm 1}] \cup \sqrt{t}\mathbf{Z}[t^{\pm 1}]$. From our Corollary 1.3 follows that if $t^{2k} = 1$, $t \neq -1$, then t_{2k} move changes $\Delta_L(t)$ by the factor $\varepsilon = t^k = \mp 1$ (i.e. $\Delta_{t_{2k}(L)}(t) = \varepsilon \Delta_L(t)$). Therefore t_{2k} moves have, more less, the same influence on $\Delta_L(t)$ as more general $t_{2k,2}$ moves; however it is not true that every $t_{2k,2}$ move is a combination of t_{2k} moves (see Example 3.8(b)).

Proof. Consider a small ball B^3 in which $t_{2n,q}$ move takes place (Fig. 2.2). $B^3 \cap L$ consists of m parallel strings.



$$m = 3$$

Fig. 2.2

$\Sigma^3 - L - \text{int} B^3$ is homeomorphic to $\Sigma^3 - t_{2n,q}(L) - \text{int} B^3$ and the fundamental groups of these spaces have the following presentation:

$$\{a_1, \dots, a_{m-1}, a'_1, \dots, a'_m, x_1, \dots : r_1, \dots\},$$

where $a_1, \dots, a_{m-1}, a'_1, \dots, a'_m$ form a basis of the free group $\pi_1(\partial B^3 - L)$; see Fig. 2.2(a).

$\Sigma^3 - L$ and $\Sigma^3 - t_{2n,q}(L)$ can be obtained from $\Sigma^3 - L - B^3$ by adding $m - 1$ two-disks in the appropriate way (Fig. 2.2(b)). Therefore

$$\pi_1(\Sigma^3 - L) = \{a_1, \dots, a_{m-1}, a'_1, \dots, a'_m, x_1, \dots : u_1, u_2, \dots, u_{m-1}, r_1, \dots\},$$

where $u_i = a'_i a_i^{-1}$, $i = 1, \dots, m - 1$ (see Fig. 2.2(b)), and

$$\pi_1(\Sigma^3 - t_{2n,q}(L)) = \{a_1, \dots, a_{m-1}, a'_1, \dots, a'_m, x_1, \dots : \tau(u_1), \tau(u_2), \dots, \tau(u_{m-1}), r_1, \dots\},$$

where $\tau(u_i) = (a'_m, \dots, a'_1)^n a'_i (a'_m, \dots, a'_1)^{-n} a_i'^{-1}$.

Consider the natural projections $p = p_2 p_1 : \pi_1(\Sigma^3 - L) \xrightarrow{P_1} H_1(\Sigma^3 - L) \xrightarrow{P_1} \mathbf{Z}$, where p sends meridians of L onto t - a generator of integers, and $p' : \pi_1(\Sigma^3 - t_{2n,q}(L)) \rightarrow \mathbf{Z}$. Then

$p(a_i) = p(a'_i) = t^{\mp 1}$ and $p(a'_m a'_{m-1} \dots a'_1) = t^q = p(a_m a_{m-1} \dots a_1)$ (without the loss of generality one can assume that the crossing number of D^2 and L is nonnegative so equal to q).

In particular if i and i' are embeddings of $\Sigma^3 - L - B^3$, in $\Sigma^3 - L$ and $\Sigma^3 - t_{2n,q}(L)$ respectively then $pi_* = p'i'_*$ (lack of this condition was the source of the mistake in the Fox paper [Fo-1]).

Now one can use Fox calculus to find Alexander-Fox modules of group representations $p : \pi_1(\Sigma^3 - L) \rightarrow \mathbf{Z}$ and $p' : \pi_1(\Sigma^3 - t_{2n,q}(L)) \rightarrow \mathbf{Z}$, and because

$$p_*\left(\frac{\partial u_i}{\partial a'_j}\right) = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases} \quad \text{and}$$

$$p'_*\left(\frac{\partial \tau(u_i)}{\partial a'_j}\right) = \begin{cases} \frac{(1-t)(1-t^{nq})t^{bj}}{1-t^q}, \text{ where } t^{bj} = p(a'_m \dots a'_{j+n}) & \text{if } i \neq j \\ \frac{(1-t)(1-t)^{nq}t^{bj}}{1-t^q} + t^{nq} & \text{if } i = j \end{cases}$$

therefore

$$p_*\left(\frac{\partial u_i}{\partial a'_j}\right) = p'_*\left(\frac{\partial \tau(u_i)}{\partial a'_j}\right) \bmod \frac{(1-t)(1-t^{nq})}{1-t^q} \text{ and one gets:}$$

Lemma 2.2. *The Alexander-Fox modules of $p : \pi_1(\Sigma^3 - L) \rightarrow \mathbf{Z}$ and $p' : \pi_1(\Sigma^3 - t_{2n,q}(L)) \rightarrow \mathbf{Z}$ can be represented by the matrices which are the same modulo $\frac{(1-t)(1-t^{nq})}{1-t^q}$.*

Theorem 2.1 follows immediately from the lemma.

□

Corollary 2.3. (a) Let qn be a multiple of s and $k = \frac{n \cdot \gcd(q,s)}{s}$; where $\gcd()$ is the greatest common divisor, then a $t_{2n,q}$ move does not change $H_1(M_L^{(s)}, \mathbf{Z}_k)$.

In particular

- (b) If q is a multiple of s (e.g. $q = 0$) then a $t_{2n,q}$ move does not change $H_1(M_L^{(s)}, \mathbf{Z}_n)$.
- (c) Let n be a multiple of s , and s and q are coprime then a $t_{2n,q}$ move does not change $H_1(M_L^{(s)}, \mathbf{Z})$.

Proof. Alexander matrices can be used to describe $H_1(M_L^{(s)}, \mathbf{Z})$ as $\mathbf{Z}[\mathbf{Z}_s]$ module ($t^s = 1$). Then we use Lemma 2.2. $\square$

Corollary 2.4. (a) Let $\bar{u}_m(L)$ denote the minimal number of $t_{2n,q}$ moves (we allow different n or q) but the number of strings involved in a $t_{2n,q}$ must be less or equal m) which are needed to change a given link L into unlink then

$$|e_s - s(c(L) - 1)| \leq (s - 1)(m - 1)\bar{u}_m(L),$$

where $c(L)$ is the number of components of L and e_s is the minimal number of generators of $H_1(M_L^{(s)}, \mathbf{Z})$. In particular for $\bar{u}(L) = \bar{u}_2(L)$ one gets:

- (b) [Kin-1] The minimal number of t_{2n} or $\bar{t}_{2n}$ moves which are needed to change a given link L into unlink ($\bar{u}(L)$), satisfies:

$$|e_s - s(c(L) - 1)| \leq (s - 1)\bar{u}(L).$$

Proof. If T_n is a trivial link of n components then $e_s(T_n) = s(n - 1)$. By the proof of Lemma 2.2, $H_1(M_{t_{2n,q}L}^{(s)}, \mathbf{Z})$ has a presentation which differs from a presentation of $H_1(M_L^{(s)}, \mathbf{Z})$ at most in $(s - 1)(m - 1)$ rows (we use additionally the fact that $\frac{t^s - 1}{t - 1} = 1 + \dots + t^{s-1}$ is an annihilator of $H_1(M_L^{(s)}, \mathbf{Z})$ (see [B-Z]), so

$$|e_s(t_{2n,q}(L)) - e_s(L)| \leq (s - 1)(m - 1).$$

$\square$

The Fox method (and Lemma 2.2) can be modified so that one can get the result about t_k moves, for k odd, analogous to Corollary 2.3 (compare [Ki]).

Consider a small ball B^3 in which a t_k move takes place (Fig. 2.3).




Fig. 2.3

$\Sigma^3 - L - \text{int} B^3$ is homeomorphic to $\Sigma^3 - t_k(L) - \text{int} B^3$ and the fundamental groups of them have the following presentation:

$$\{a, b, c, x_1, x_2, \dots : r_1, r_2\},$$

where a , b and c are classes of curves (generators of $\pi_1(\partial B^3 - L)$) shown on Fig. 2.4.




Fig. 2.4

If we add a 2-handle along $u = ac^{-1}$ we get $\Sigma^3 - L$ and if we add a 2-handle along $(ba)^{(k-1)/2}b(ba)^{-(k-1)/2}c^{-1}$ we get $\Sigma^3 - t_k(L)$; therefore

$$\pi_1(\Sigma^3 - L) = \{a, b, c, x_1, x_2, \dots : ac^{-1} = 1, r_1, r_2, \dots\},$$

$$\pi_1(\Sigma^3 - t_k(L)) = \{a, b, c, x_1, x_2, \dots : (ba)^{(k-1)/2}b(ba)^{-(k-1)/2}c^{-1}, r_1, r_2, \dots\}.$$

Consider the natural projections $p : \pi_1(\Sigma^3 - L) \rightarrow \mathbf{Z}$ and $p' : \pi_1(\Sigma^3 - t_k(L)) \rightarrow \mathbf{Z}$. Then we have:

$$p(a) = p(b) = p(c) = t, \quad p'(a) = p'(b) = p'(c) = t.$$

Now we calculate that $(r_0 = (ba)^{(k-1)/2}b(ba)^{-(k-1)/2}c^{-1})$:

$$p\left(\frac{\partial ac^{-1}}{\partial a}\right) = 1, \quad p'\left(\frac{\partial r_0}{\partial a}\right) = 1 - \frac{t^k + 1}{t + 1},$$

$$p\left(\frac{\partial ac^{-1}}{\partial b}\right) = 0, \quad p'\left(\frac{\partial r_0}{\partial b}\right) = \frac{t^k + 1}{t + 1},$$

$$p\left(\frac{\partial ac^{-1}}{\partial c}\right) = -1, \quad p'\left(\frac{\partial r_0}{\partial c}\right) = -1,$$

and we get:

Lemma 2.5. *The Alexander-Fox modules of L and $t_k(L)$ can be presented by the following matrices.*

L	a	b	c	x_1	x_2	$\dots$	$t_k(L)$	a	b	c	x_1	x_2	$\dots$
$ac^{-1} = 1$	1	0	-1	0	0	$\dots$	r_0	$1 - \frac{t^k+1}{t+1}$	$\frac{t^k+1}{t+1}$	-1	0	0	$\dots$
r_1	*	*		*	*		r_1	*	*		*	*	*
$\dots$			$\dots$				$\dots$			$\dots$			

Corollary 2.6. (a) *For $\frac{t^k+1}{t+1} = 0$ (k -odd) t_k -equivalent links have the same Alexander module, in particular*

$$(b) \Delta_{t_k(L)}(t) \equiv \mp t^i \Delta_L(t) \pmod{\frac{t^k+1}{t+1}}.$$

In fact from Corollary 1.3 follows that for a normalized Alexander polynomial $\Delta_L(t) \equiv \mp i \Delta_{t_k(L)}(t) \pmod{\frac{t^k+1}{t+1}}$ or precisely

$$\Delta_L(t) \equiv t^{k/2} \Delta_{t_k(L)}(t) \pmod{\frac{t^k+1}{t+1}}.$$

We can slightly generalize the results of Wendt and Kinoshita using Lemma 2.5.

Corollary 2.7. *Let $\bar{\bar{u}}_n(L)$ denote the minimal number of $\bar{t}_{2k}$ or t_k moves which are needed to change a given oriented link L into unlink of n components, then*

$$|e_s - s(n-1)| \leq (s-1) \bar{\bar{u}}_n(L),$$

where e_s is the minimal number of generators of $H_1(M_L^{(s)}, \mathbf{Z})$.

3. APPLICATIONS AND SPECULATIONS

We start this part by proving two "folklore" results which link Goeritz and Seifert matrices with t_k or $\bar{t}_k$ moves.

Theorem 3.1. (a) *There exist Goeritz's matrices for L and $t_k(L)$ (or $\bar{t}_k(L)$) which are the same modulo k .*

(b) *t_k and $\bar{t}_k$ -moves preserves $H_1(M_L^{(2)}, \mathbf{Z}_k)$.*

Proof. For the convenience we start from the definition of Goeritz's matrix ([Goe, Gor]). Colour the regions of the diagram of an unoriented link alternately black and white, the unbounded region X_0 being coloured white, and number the other white

regions $X_1, \dots, X_n$. Assign an incidence number $\eta(p) = \mp 1$ to each crossing point p as shown in Fig. 3.1. Then define $n \times n$ Goeritz's matrix $G = (g_{ij})$ by




Fig. 3.1

$$g_{ij} = \begin{cases} \sum \eta(c) & \text{summed over crossings points } p \text{ adjacent to } X_i \text{ and } X_j \\ & \text{if } i \neq j \text{ (} i \cdot j \geq 1 \text{)} \\ -\sum \eta(c) & \text{summed over crossings points } p \text{ adjacent to} \\ & X_i \text{ and to some } X_j \text{ (} i \neq j \text{) if } i = j \text{ (} i \geq 1 \text{).} \end{cases}$$

Now consider the Fig. 3.2 with white regions X_i and X_j .




Fig. 3.2

There are two possible cases:

- (i) $X_i = X_j$, then $G_L = G_{L'}$; in fact L is isotopic to L' .
- (ii) $X_i \neq X_j$, we can assume that $i = 0$ and $j = 1$ then

$$G_{L'} = \begin{bmatrix} g_{11} + k, & g_{12} & \cdots, & g_{1n} \\ & \cdots & & \\ g_{n1}, & g_{n2} & \cdots, & g_{nn} \end{bmatrix} \text{ where } G_L = (g_{ij})$$

.

The part (b) of the Theorem 3.1 follows from the fact that G_L is a presentation matrix for $H_1(M_L^{(2)}, \mathbf{Z})$. $\square$

An alternative proof of (b) can be given by considering Dehn surgery on $M_L^{(2)}$ corresponding to t_k or $\bar{t}_k$ move on L .

Theorem 3.2. (a) Consider a $t_{2k,0}$ move of Fox (e.g. $\bar{t}_{2k}$ move), then there exist Seifert matrices for L and $t_{2k,0}(L)$ which are the same modulo k .
 (b) $t_{2k,0}$ move preserves $H_1(M_L^{(s)}, \mathbf{Z}_k)$ for any s .

Proof. One can find a Seifert surface S for L which cuts the disk D^2 which supports the $t_{2k,0}$ move, as shown in Fig. 3.3. Then the Seifert matrix for L defined by S and for $t_{2k,0}(L)$ defined by $t_{2k,0}(S)$ satisfy the condition (a).




Fig. 3.3

(b) follows from (a) because a presentation matrix for $H_1(M_L^{(s)}, \mathbf{Z})$ can be built of blocks of the shape $\mp V$, $\mp V^T$, $\mp(V + V^T)$, where V is a Seifert matrix of L and V^T its transpose. On the other hand, (b) is a special case of Theorem 2.3(b). $\square$

Example 3.3. (a) *The trivial knot (T_1) and the (right handed) trefoil knot (3_1) are t_4 equivalent. The figure eight knot (4_1) and the 5_2 knot are t_4 equivalent however they are not t_4 equivalent to T_1 or 3_1 .*
 (b) *T_1 and 5_2 are $\bar{t}_4$ equivalent. 3_1 and 4_1 are $\bar{t}_4$ equivalent but they are not $\bar{t}_4$ equivalent to T_1 or 5_2 .*

First parts of (a) and (b) are illustrated in Fig. 3.4.




Fig. 3.4

The second parts follow from Corollaries 1.2 and 1.8 (t_4 move changes $P_L(a, \sqrt{2})$ by the factor $-a^{-4}$ and $\bar{t}_4$ preserves $P_L(1, z)$) and the following computation:

$$P_{T_1}(a, z) = 1$$

$$P_{3_1}(a, z) = -a^{-4} - 2a^{-2} + z^2a^{-2}; \quad P_{3_1}(a, \sqrt{2}) = -a^{-4}, \quad P_{3_1}(1, z) = z^2 - 3$$

$$P_{4_1}(a, z) = -a^{-2} - 1 - a^2 + z^2; \quad P_{4_1}(a, \sqrt{2}) = -a^{-2} + 1 - a^2; \quad P_{4_1}(1, z) = z^2 - 3$$

$$P_{5_2}(a, z) = -a^{-2} + a^4 + a^6 + z^2(a^2 - a^4); \quad P_{5_2}(a, \sqrt{2}) = -a^4(-a^{-2} + 1 - a^2); \quad P_{5_2}(1, z) = 1.$$

Example 3.4. Every closed 3-braid knot is t_4 equivalent to the trivial knot or the figure eight knot. It is not an unexpected result because the quotient group $B_3/(\delta_1^4)$ is finite [Cox]. In fact a calculation shows that $B_3/(\delta_1^4)$ has only two classes of

knots (represented by T_1 and 4_1). Because all presentations of 4_1 as a 3-braid (e.g. $\delta_1\delta_2^{-1}\delta_1\delta_2^{-1}$) have the same exponent sum (equal to 0) therefore for every knot K which is t_4 equivalent to 4_1 , each of its presentation as a 3-braid has the same exponent sum (equal to $4|4_1, K|_{t_4}^{\text{lev}}$; compare Corollary 1.2). More in this direction can be got using other t_k moves, compare Example 3.11, however it has been generally proved by H.Morton [Mo] and J.Birman that if L is not a $(2, k)$ torus link then the exponent sum of L does not depend on the presentation of L as a closed 3-braid.

Example 3.5. Consider the following theorem of H.Murakami [Mur-1] (see also [L-M-2]):

$$P_L(1, \sqrt{2}) = V_L(i) = \begin{cases} (\sqrt{2})^{c(L)-1}(-1)^{\text{Arf}(L)} & \text{if Arf}(L) \text{ exists} \\ 0 & \text{otherwise ,} \end{cases}$$

where $c(L)$ denotes the number of components of L , $\text{Arf}(L)$ is the Arf (or Robertello) invariant (see [Rob] or [Ka-2]), and $t = i$ in $V_L(t)$ should be understood as $t^{1/2} = -e^{\pi i/4}$. Notice that our convention differs slightly from that of [L-M-1] or [L-M-2] namely $P_L(a, z) = P_L(\ell, -m) = (-1)^{c(L)-1}P_L(\ell, m)$.

It follows from Corollaries 1.2 and 1.8 that t_4 move changes $P_L(1, \sqrt{2})$ by factor -1 and $\bar{t}_4$ move preserves $P_L(1, \sqrt{2})$. Furthermore for T_n - the trivial link of n components $P_{T_n}(1, \sqrt{2}) = (\sqrt{2})^{n-1}$. On the other hand the Arf invariant of a trivial link is equal to zero, t_4 move changes the Arf invariant (if defined) and $\bar{t}_4$ move preserves it (see [Ka-2]). Therefore the Murakami theorem follows immediately from the above observations for a link which is $t_4, \bar{t}_4$ equivalent to a trivial link (i.e. a link which can be obtained from a trivial one using t_4 and $\bar{t}_4$ moves). This should be confronted with the following conjecture:

Conjecture 3.6. (*Kawauchi - Nakanishi*)

- (a) *If two links L_1 and L_2 are homotopic then they are $t_4, \bar{t}_4$ equivalent¹. In particular:*
- (b) *Every knot is $t_4, \bar{t}_4$ equivalent to the unknot.*

Conjecture 3.6 has been verified for the 2-bridge links, closed 3-string braids and pretzel links.

Example 3.7. *Consider the following t_{Δ^2} -move (Δ^2 -twist) on oriented diagrams of links (Fig. 3.5).*




Fig. 3.5

¹Added for e-print: The conjecture has been disproved in [D-P-2] for links of three or more components. For two component links it is still an open problem whether any such link is $t_4, \bar{t}_4$ equivalent to T_2 or the Hopf link.

A t_4 move can be obtained from a t_{Δ^2} -move (and isotopy) as it is illustrated in Fig.3.6.




Fig. 3.6

J. Birman and B. Wajnryb [B-W] have proven that two links are t_{Δ^2} equivalent iff they have the same number of components and the same number of components with odd linking number with the rest of the link. Because a $\bar{t}_4$ move preserves the number of components and all linking numbers modulo 2, therefore it can be obtained as a combination of $t_{\Delta^2}^{\mp 1}$ moves. In fact it follows from [B-W] that in order to get $\bar{t}_4$ move we can always use an even number of $t_{\Delta^2}^{\mp 1}$ moves. Furthermore a t_{Δ^2} move changes the Arf invariant (if it exists) and therefore $V_L(i) = -V_{t_{\Delta^2}(L)}(i)$. The last equality can be also proven elementary without using [B-W]. Finally observe that not every t_{Δ^2} move is a combination of t_4 , $\bar{t}_4$ moves. The reason is that t_4 and $\bar{t}_4$ moves

preserve all linking numbers mod 2 but it is not always the case for a t_{Δ^2} move (see Fig. 3.7 for an example of links which are t_{Δ^2} equivalent but not $t_4, \bar{t}_4$ equivalent).




Fig. 3.7

Example 3.8. (a) a t_{Δ^2} move is a special case of $t_{2,3}$ moves of Fox but it follows from [B-W] that any $t_{2,3}$ move is a combination of t_{Δ^2} moves. In fact, every $t_{2,3}$ move preserves the number of components and the number of components with odd linking number with the rest of the link. Similarly any $t_{2,2q+1}$ is a combination of t_{Δ^2} moves.

- (b) a t_4 move is a special case of $t_{4,2}$ moves of Fox. There are $t_{4,2}$ equivalent links which are not t_4 equivalent. (Fig. 3.8).




Fig. 3.8

Two links of Fig. 3.8 are not t_4 equivalent because their sublinks of Fig. 3.9 are not t_4 equivalent.




Fig. 3.9

Namely $P_{T_2}(a, \sqrt{2}) = \frac{a+a^{-1}}{\sqrt{2}}$ and $P_K(a, \sqrt{2}) = \frac{a^{-5}-a^{-3}+2a^{-1}}{\sqrt{2}}$, therefore by Corollary 1.2 T_2 and K are not t_4 equivalent.

- Example 3.9.** (a) The square knot $(3_1 \# \bar{3}_1)$, the (right-handed) granny knot $(3_1 \# 3_1)$, and T_3 (the trivial 3-component link) are t_3 equivalent.
- (b) The trefoil knot (3_1) and T_2 (the trivial link of 2-components) are t_3 equivalent.
- (c) The knots 5_2 , 6_3 , the Hopf link (2_1^2) , the Borromean rings (6_2^3) and the unknot (T_1) are t_3 equivalent.
- (d) The figure eight knot (4_1) and the knot 9_{42} (in the Rolfsen notation [Rol]) are t_3 equivalent.
- (e) No links from different classes ((a), (b), (c), (d)) are t_3 equivalent however links of (c) and (d) are t_3 , $\bar{t}_3$ equivalent (i.e. there is a sequence of $t_3^{\pm 1}$ or $\bar{t}_3^{\pm 1}$ moves which lead from one link to another) and there is no more t_3 , $\bar{t}_3$ equivalences among the above links.

The t_3 and t_3 , $\bar{t}_3$ equivalences are illustrated in Fig. 3.10, 3.11, 3.12 and 3.13.




Fig. 3.10




Fig. 3.11




Fig. 3.12




Fig. 3.13

The first part of (e) follows Corollary 1.2 (a t_3 move changes $P_L(a, 1)$ by the factor $-a^{-3}$) and the following computation:

$$P_{T_1}(a, 1) = 1, \quad P_{T_2}(a, 1) = a + a^{-1}, \quad P_{T_3}(a, 1) = a^{-2} + 2 + a^2, \quad P_{4_1}(a, 1) = -a^{-2} - a^2.$$

The last statement of (e) follows from the fact that different trivial links are not $t_3, \bar{t}_3$ equivalent (see Lemma 3.10(c) below).

Lemma 3.10. Consider the Jones polynomial $V_L(t)$ for $t = e^{\pi i/3}$ ($t^{1/2} = -e^{\pi i/6}$), then

$$(a) \quad V_{t_3(L)}(e^{\pi i/3}) = iV_L(t)$$

$$(b) \quad V_{\bar{t}_3(L)}(e^{\pi i/3}) = \begin{cases} (-1)^\lambda iV_L(t) & \text{if two components of } L \text{ are involved in } \bar{t}_3 \text{ move} \\ (-1)^\lambda V_L(t) & \text{if one component of } L \text{ is involved in } \bar{t}_3 \text{ move} \end{cases}$$

λ depends on the linking numbers of components of L and $\bar{t}_3(L)$ and on an orientation of $\bar{t}_3(L)$ (see Theorem 1.13).

- (c) The trivial links T_k and T_j ($k \neq j$) are not $t_3, \bar{t}_3$ equivalent and $V_{T_k}(e^{\pi i/3}) = (\sqrt{3})^{k-1}$.

Proof. (a) follows from Corollary 1.4, and (b) from Theorem 1.13. (c) follows from (a) and (b). $\square$

Example 3.11. Every closed 3-braid link is t_3 equivalent to T_1, T_2, T_3 or the figure eight knot (in fact one can go from any closed 3-braid link to one of these links using t_3 moves and regular isotopy). Because all presentations of 4_1 and T_3 as closed 3-braids have the same exponent sum (equal to 0) therefore for any link L which is t_3 equivalent to 4_1 or T_3 , each of its presentation as a 3-braid has the same exponent sum (equal to $3|4_1, L|_{t_3}^{\text{lev}}$ or $3|T_3, L|_{t_3}^{\text{lev}}$). Consider, for example, the closed 3-braid knot $\delta_1^4 \delta_2^{-1} \delta_1 \delta_2^{-4}$ (Fig. 3.14). It is t_3 equivalent to the figure eight knot so now we know that all presentations of this knot as a 3-braid have the exponent sum equal to zero; on the other hand, the knot is t_4 equivalent to the unknot so the method of Example 3.4 would not suffice to get the unique exponent sum.




Fig. 3.14

Example 3.12. Consider the following theorem of W.B.R.Lickorish and K.Millett [L-M-2] (conjectured by J.Birman and partially proved by V.Jones).

$$P_L(e^{\pi i/6}, 1) = V_L((e^{\pi i/3}) = \mp i^{c(L)-1} (i\sqrt{3})^{\text{Dim } H_1(M_L^{(2)}, \mathbf{Z}_3)},$$

where $c(L)$ denotes the number of components of L and $t^{1/2} = e^{-\pi i/6}$ in $V_L(t)$.

It follows from Lemma 3.10 that t_3 and $\bar{t}_3$ moves change $V_L((e^{\pi i/3})$ by factors ∓ 1 or $\mp i$ and the second case happens if the move changes the number of components. On the other hand t_3 and $\bar{t}_3$ moves preserve $H_1(M_L^{(2)}, \mathbf{Z}_3)$ (Theorem 3.1(b)) and for the trivial link T_n , $\text{Dim } H_1(M_L^{(2)}, \mathbf{Z}_3) = n - 1$. Therefore the formula of Lickorish-Millett holds immediately from the above observations for a link which is t_3 , $\bar{t}_3$ equivalent to a trivial link (the sign in formula can be found using Lemma 3.10; it was identified generally by A.Lipson [Lip]). This should be confronted with the following conjecture.

Conjecture 3.13. (Montesinos-Nakanishi). Every link is t_3 , $\bar{t}_3$ equivalent to a trivial link.²

It is an easy (but tedious) task to check the conjecture for closed n -braids ($n \leq 5$) and n -bridge links ($n \leq 3$) because for the braid group B_n ($n \leq 5$) the group $B_n/(\delta_1^3)$ is finite ([Cox]), however the author did it only for closed 3-braids and 2-bridge links.³

Example 3.14. Consider the following theorem of Lickorish and Millett [L-M-2] and H. Murakami [Mur-2]:

²Added for e-print: The conjecture has been disproved in [D-P-1]. The smallest known counter-example has 20 crossings.

³Added for e-print: The conjecture holds for 4-bridge links [P-Ts, Tsu]. Furthermore every closed 5-braid is t_3 , $\bar{t}_3$ equivalent to a trivial link or to the closure of the 5-string braid $(\delta_1\delta_2\delta_3\delta_4)^{10}$ [Chen]. The last link is a counter-example to Montesinos-Nakanishi conjecture [D-P-1].

$$P_L(1, 1) = (2)^{(1/2)\text{Dim } H_1(M_L^{(3)}, \mathbf{Z}_2)}.$$

It follows from Corollaries 1.2 and 1.8 that t_3 and $\bar{t}_4$ moves preserve $P_L(1, 1)$. Furthermore, $P_{T_n}(1, 1) = 2^{n-1}$. On the other hand $\text{Dim } H_1(M_{T_n}^{(3)}, \mathbf{Z}_2) = 2(n-1)$ and $\bar{t}_4$ moves preserve $H_1(M_L^{(3)}, \mathbf{Z}_2)$. It can be shown, using the Fox approach that t_3 -moves preserve $H_1(M_L^{(3)}, \mathbf{Z}_2)$ ([P-3]). Therefore the formula of Lickorish-Millett-Murakami follows immediately from the above observations for a link which is $t_3, \bar{t}_4$ equivalent to a trivial link (i.e. a link which can be got from a trivial one using t_3 and $\bar{t}_4$ moves and isotopy). This leads to the following conjecture.

Conjecture 3.15. *Every link is $t_3, \bar{t}_4$ equivalent to a trivial link.*

The author has checked the conjecture for closed 3-braid links (see the remark after Conjecture 3.13).⁴

For t_5 and $\bar{t}_4$ moves the analogy of Conjecture 3.15 does not hold. For example, trivial links, the trefoil knot, 8_5 knot and 8_{18} knot ([Rol], see Fig. 3.15) are not pairwise $t_5, \bar{t}_4$ equivalent. The reason is that by Corollary 1.8 a $\bar{t}_4$ move does not change $P_L(1, z)$ and by Corollary 1.2 t_5 move changes $P_L(1, \frac{1+\sqrt{5}}{2})$ by the factor -1 (notice that $2 \cos(\pi i/5) = \frac{1+\sqrt{5}}{2}$); on the other hand all mentioned above links have pairwise different absolute values of $P_L(1, \frac{1+\sqrt{5}}{2})$:

$$P_{T_n}(1, \frac{1+\sqrt{5}}{2}) = (\sqrt{5} - 1)^{n-1}$$

$$P_{3_1}(1, \frac{1+\sqrt{5}}{2}) = \frac{-3+\sqrt{5}}{2}, \quad P_{8_5}(1, \frac{1+\sqrt{5}}{2}) = -4 + \sqrt{5},$$

$$P_{8_{18}}(1, \frac{1+\sqrt{5}}{2}) = \frac{1-2\sqrt{5}}{2}.$$

⁴Added for e-print: It has been checked for closed 4-braid links [Chen].



Fig. 3.15

There is no chance for anything analogous to Conjectures 3.6 or 3.13 for $t_k, \bar{t}_k$ moves, $k \geq 5$ (i.e. that all links are $t_k, \bar{t}_k$ equivalent to the trivial links). In particular V.Jones ([Jo-3]; Corollary 14.7) proved that the set $\{|V_L(e^{\pi i/5})| : L \text{ is a link}\}$ is dense in $\langle 0, \infty \rangle$. On the other hand t_5 and $\bar{t}_5$ moves do not change the absolute value of $V_L(e^{\pi i/5})$ (see Corollary 1.4 and Theorem 1.13), and for trivial links, the values $|V_{T_n}(e^{\pi i/5})| = (2 \cos \pi/10)^{n-1}$ are discrete in $\langle 0, \infty \rangle$.

There are natural relations between t_k moves and signatures of links; we will list here some examples of such relations. For convenience, we start from the definition of the Tristram-Levine signature (see [Gor, P-T-2] or [P-1]). Let A_L be a Seifert matrix of a link L . For each complex number ξ ($\xi \neq 1$) consider Hermitian matrix $A_L(\xi) = (1 - \bar{\xi})A_L + (1 - \xi)A_L^T$. The signature of this matrix, $\sigma_L(\xi)$ is called the Tristram-Levine signature of the link L . The classical signature σ satisfies $\sigma_L = \sigma_L(0)$.

Theorem 3.16. (a) For any t_k move on an oriented link L

$$k - 2 \leq \sigma_L - \sigma_{t_k(L)} \leq k,$$

(b) $0 \leq \sigma_{\bar{t}_{2k}(L)} - \sigma_L(\xi) \leq 2$ if $\operatorname{Re}(1 - \xi) \geq 0$,

(c) $\sigma_L(\xi_0) - \sigma_{t_4}(\xi_0) = 2$ if $P(i, \sqrt{2}) \neq 0$ and $\xi_0 = 1 - e^{\pi i/4} = \frac{2-\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}$,

- (d) If two links L_1 and L_2 are t_4 equivalent then $|L_1, L_2|_{t_4}^{\text{lev}} = (1/2)(\sigma_{L_1}(\xi_0) - \sigma_{L_2}(\xi_0))$, provided $P_{L_1}(i, \sqrt{2}) \neq 0$.

Proof. (a) We use the formula of C.McA.Gordon, R.A.Litherland [G-L] and A.Marin, which links the signature of Goeritz matrix of a link with a classical signature. We use the same notation as in the proof of Theorem 3.1. Divide the crossings of a given oriented link L into two types as shown in Fig. 3.16.




Fig. 3.16

Define $\mu = \sum \eta(p)$, summed over all crossing points of type II then $\sigma_L = \sigma(G_L) - \mu(L)$. Furthermore we have $\mu(t_k(L)) - \mu(L) = k$ (see Fig. 3.2), and from the form of the matrices $G_{L'} = G_{t_k(L)}$ and G_L (see proof of Theorem 3.1) follows that $-2 \leq \sigma(G_L) - \sigma(G_{t_k(L)}) \leq 0$ and therefore $-2 \leq \sigma_L + \mu(L) - \sigma_{t_k(L)} - \mu(t_k(L)) \leq 0$ and Theorem 3.16(a) follows.

To prove (b), we have to choose a proper Seifert surface from which we will find the adequate Seifert matrix so one could easily compare the Levine-Tristram signature

for L and $\bar{t}_{2k}(L)$. We can assume that Seifert surfaces for L and $\bar{t}_{2k}(L)$ looks locally as on Fig. 3.17 (or $\bar{t}_{2k}(L)$ is isotopic to L).




Fig. 3.17

Then the Seifert matrices (in appropriate basis) are of the form :

$$A_{\bar{t}_{2k}(L)} = \begin{bmatrix} A_L & \alpha \\ \beta & q+k \end{bmatrix},$$

$$A_L = \begin{bmatrix} A_L & \alpha \\ \beta & q \end{bmatrix},$$

where A_L is the Seifert matrix of L , α is a column, β is a row and q is a number (compare [Ka-1, P-T-2] or [P-1]). Therefore

$$A_{\bar{t}_{2k}(L)}(\xi) = \begin{bmatrix} A_L(\xi) & a \\ a^{-T} & m+k(2-\xi-\bar{\xi}) \end{bmatrix},$$

$$A_L(\xi) = \begin{bmatrix} A_L(\xi) & a \\ a^{-T} & m \end{bmatrix},$$

where $a = (1-\bar{\xi})\alpha + (1-\xi)\beta^T$ and $m = ((1-\bar{\xi}) + (1-\xi))q$. Because $2-\xi-\bar{\xi} \geq 0$, so $0 \leq \sigma(A_{\bar{t}_{2k}(L)}(\xi)) - \sigma(A_L(\xi)) \leq 2$ and the proof of (b) is finished.

To prove (c) we need further characterization of the Tristram-Levine signature, given in [P-T-2] (see also [P-1]); Assume $|1-\xi| = 1$, we have :

- (i) $\text{Det } iA_L(\xi) = P_L(i, 2-\xi-\bar{\xi}) = \Delta_L(t')$ (for $\sqrt{t'} = -i(1-\xi)$),

- (ii) $i\sigma_L(\xi) = \frac{\Delta_L(t')}{|\Delta_L(t')|}$ if $\Delta_L(t') \neq 0$,
 (iii) $0 \leq \sigma_L(\xi) - \sigma_{t_4(L)}(\xi) \leq 4$ if $\operatorname{Re}(1 - \xi) \geq 0$.

((iii) can be got using (b) two times with $k = 1$; $\bar{t}_2$ moves are equivalent to t_2 moves).

Now consider the case when $2 - \xi_0 - \bar{\xi}_0 = \sqrt{2}$ ($1 - \xi_0 = e^{\pi i/4}$). Then by Corollary 1.3, $\Delta_{t_4(L)}(t') = -\Delta_L(t')$. Therefore by (ii) and (iii) $\delta_L(\xi_0) - \delta_{t_4(L)}(\xi_0) = 2$. (d) follows immediately from (c). $\square$

One can expect interesting relations between t_k moves and non-cyclic coverings of links. We limit ourself to two examples, first of which was suggested by R.Campbell.

Example 3.17. (a) *A link diagram is 3-coloured if every overpass is coloured, say, red, yellow or blue, at least two colours are used and at any given crossing either all three colours appear or only one colour appears [Fo-2]. Then if a link L_1 is $t_3, \bar{t}_3$ equivalent to L_2 then either both links are 3-coloured or none of them are 3-coloured. In particular a link which is $t_3, \bar{t}_3$ equivalent to a trivial link of more than one component is 3-coloured. The proof is illustrated in Fig. 3.18. The link 6_3^2 [Rol] is 3-coloured in Fig. 3.19.*




Fig. 3.18

- (b) 3-colouring corresponds to an epimorphism $\pi_1(S^3 - L) \rightarrow S_3$; more generally we have: If a knot K_1 is $t_{2p}, \bar{t}_{2p}$ equivalent to K_2 (p - prime) then either both knots or none of them have dihedral representations i.e. epimorphism

$$\pi_1(S^3 - K) \rightarrow D_{2p} = \{a, b : a^2 = 1, b^p = 1, aba = b^{-1}\}.$$

It follows from the fact that $t_{2p}, \bar{t}_{2p}$ moves preserve $H_1(M_K^{(2)}, \mathbf{Z}_p)$ (Theorem 3.1(b)) and from the result of Fox that the epimorphism exists iff $H_1(M_K^{(2)}, \mathbf{Z}_p)$, is nontrivial [Fo-2] (see also [B-Z]; 14.8).⁵




Fig. 3.19

REFERENCES

- [A] J.W.Alexander, *Topological invariants of knots and links*, Trans. Amer. Math. Soc., 30 (1928), 275-306.
- [Bi] J.S.Birman, *Braids, links and mapping class groups*, Ann. Math. Studies 82, Princeton Univ. Press, 1974.
- [B-W] J.S.Birman, B.Wajnryb, *Markov classes in certain finite quotients of Artin's braid group*, Israel J. Math. (to appear). Added for e-print: 56(2), 1986, 160-178.

⁵Added for e-print: These ideas have been developed in [P-5].

- [B-Z] G.Burde, H.Zieschang, *Knots*, De Gruyter Studies in Math. 5, Berlin, New York 1985.
- [Chen] Added for e-print: Q. Chen, The 3-move conjecture for 5-braids, Knots in Hellas' 98; The Proceedings of the International Conference on Knot Theory and its Ramifications; Volume 1. In the Series on Knots and Everything, Vol. 24, September 2000, pp. 36-47.
- [Con] J.H.Conway, *An enumeration of knots and links, and some of their algebraic properties*, Computational problems in abstract algebra (J.Leech, ed.), Pergamon, Oxford and New York (1969), 329-358.
- [Cox] H.S.M.Coxeter, *Factor groups of the braid group*, Proc. Fourth Canadian Math. Congress, Banff, 1957, 95-122.
- [D-P-1] Added for e-print: M. K. Dąbkowski, J. H. Przytycki, Burnside obstructions to the Montesinos-Nakanishi 3-move conjecture, i *Geometry and Topology (G&T)*, 6, June, 2002, 335-360;
e-print: <http://front.math.ucdavis.edu/math.GT/0205040>
- [D-P-2] Added for e-print: M. K. Dąbkowski, J. H. Przytycki, Unexpected connection between knot theory and Burnside groups, *Proc. Nat. Acad. Science*, 101(50), December, 2004, 17357-17360;
e-print: <http://front.math.ucdavis.edu/math.GT/0309140>
- [Fo-1] R.H.Fox, *Congruence classes of knots*, Osaka Math. J. 10 (1958), 37-41.
- [Fo-2] R.H.Fox, *Metacyclic invariants of knots and links*, Canadian J. Math. XXII (2) (1970), 193-201.
- [F-Y-H-L-M-O] P.Freyd, D.Yetter, J.Hoste, W.B.R.Lickorish, K.Millet, A.Ocneau, *A new polynomial invariant of knots and links*, Bull. Amer. Math. Soc. 12(2) (1985), 239-249.
- [Goe] L.Goeritz, *Knoten und quadratische Formen*, Math. Z. 36 (1933), 647-654.
- [Gor] C.McA.Gordon, *Some aspects of classical knot theory*, In: Knot theory, L.N.M. 685 (1978), 1-60.
- [G-L] C.McA.Gordon, R.A.Litherland, *On the signature of a link*, Inv. Math. 47 (1978), 53-69.
- [Jo-1] V.F.R.Jones, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. 12(1) (1985), 103-111.
- [Jo-2] V.F.R.Jones, *A new knot polynomial and von Neumann algebras*, Notices AMS 33(2) (1986), 219-225.
- [Jo-3] V.F.R.Jones, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (to appear). Added for e-print: 126(2), 1987, 335-388.
- [Kan] T.Kanenobu, *Examples on polynomial invariants of knots and links*, Math. Ann. 275 (1986), 555-572.
- [Ka-1] L.H.Kauffman, *The Conway polynomial*, Topology 20 (1980), 101-108.
- [Ka-2] L.H.Kauffman, *Knots*, Lecture notes, Zaragoza, Spring 1984. Added for e-print: *On knots*, Annals of Math. Studies, 115, Princeton University Press, 1987.

- [Ka-3] L.H.Kauffman, *An invariant of regular isotopy*, preprint 1985. Added for e-print: *Trans. Amer. Math. Soc.*, 318(2), 1990, 417–471.
- [Ka-4] L.H.Kauffman, *State models for knot polynomials*, preprint 1985. Added for e-print: *Topology* 26, 1987, 395–407.
- [Ki] M.E.Kidwell, *Relations between the Alexander polynomial and summit power of a closed braid*, Math. Sem. Notes Kobe 10 (1982) 387–409.
- [Kin-1] S.Kinoshita, *On Wendt's theorem of knots, I*, Osaka Math. J. 9(1) (1957) 61–66.
- [Kin-2] S.Kinoshita, *On Wendt's theorem of knots, II*, Osaka Math. J. 10 (1958) 259–261.
- [Kin-3] S.Kinoshita, *On the distribution of Alexander polynomials of alternating knots and links*, Proc. Amer. Math. Soc. 79(4) (1980), 644–648.
- [Li] W.B.R.Lickorish, *A relationship between link polynomials*, Math. Proc. Cambridge Phil. Soc. (1986), 100, 109–112.
- [L-M-1] W.B.R.Lickorish, K.C.Millet, *A polynomial invariant of oriented links*, Topology 26(1) (1987), 107–141.
- [L-M-2] W.B.R.Lickorish, K.C.Millet, *Some evaluations of link polynomials*, Com. Math. Helvetici 61(3) (1986), 349–359.
- [L-M-3] W.B.R.Lickorish, K.C.Millet, *The reversing result for the Jones polynomial*, Pacific J. Math. (to appear). Added for e-print: 124 (1986), 173–176.
- [Lip] A.S.Lipson *An evaluation of a link polynomial*, Math. Proc. Camb. Phil. Soc. 100 (1986), 361–364.
- [Mon] J.M.Montesinos *Lectures on 3-fold simple coverings and 3-manifolds*, Amer. Math. Soc. Contemp. Math. 44 (1985), 157–177.
- [Mo] H.R.Morton Personal conversation, (July 1986).
- [Mu] K.Murasugi, *On closed 3-braids*, Memoirs AMS 151 (1974) Amer. Math. Soc. Providence, R.I.
- [Mur-1] H.Murakami, *A recursive calculation of the Arf invariant of a link*, J. Math. Soc. Japan 38(2) (1986), 335–338.
- [Mur-2] H.Murakami, *Unknotting number and polynomial invariants of a link*, preprint 1985.
- [N] Y.Nakanishi, *Fox's congruence classes and Conway's potential functions of knots and links*, preprint 1986.
- [N-S] Y.Nakanishi, S.Suzuki, *On Fox's congruence classes of knots*, Osaka J. M. 24 (1987) (to appear). Added for e-print: pages 217–225.
- [P-1] J.H.Przytycki, *Survey on recent invariants in classical knot theory*, preprint, Warsaw University, 1986. Added for e-print: Part of: *Teoria węzłów: podejście kombinatoryczne*, (Knots: combinatorial approach to the knot theory), Script, Warsaw, August 1995, 240+XLVIIIpp. (Extended version for Cambridge University Press, to appear).

- [P-2] J.H.Przytycki, t_k equivalence of links and Conway formulas for the Jones-Conway and Kauffman polynomials, preprint 1986. Added for e-print: *Bull. Polish Acad. Sci. Math.*, 36(11-12), 1988, 675-680.
- [P-3] J.H.Przytycki, *Plans' theorem for links: An application of t_k moves*, Bull. Canad. Math. Soc., (to appear). Added for e-print: 31(3), 1988, 325-327.
- [P-4] J.H.Przytycki, *On Murasugi and Traczyk criteria for periodic links*, preprint, March 1987. Added for e-print: *Math. Ann.*, 283, 1989, 465 - 478.
- [P-5] Added for e-print: J. H. Przytycki, 3-coloring and other elementary invariants of knots, Banach Center Publications, Vol. 42, *Knot Theory*, 1998, 275-295.
- [P-T-1] J.H.Przytycki, P.Traczyk, *Invariants of links of Conway type*, Kobe J. Math. (to appear). Added for e-print: 4, 1987, 115-139.
- [P-T-2] J.H.Przytycki, P.Traczyk, *Conway algebras and skein equivalence of links*, preprint, 1985. Added for e-print: Part of the paper published in *Proc. Amer. Math. Soc.*, 100(4), 1987, 744-748.
- [P-Ts] Added for e-print: J. H. Przytycki, T.Tsukamoto, The fourth skein module and the Montesinos-Nakanishi conjecture for 3-algebraic links, *J. Knot Theory Ramifications*, 10(7), 2001, 959-982.
<http://front.math.ucdavis.edu/math.GT/0010282>
- [Re] K.Reidemeister, *Knotentheorie*, Ergebn, Math. Grenzgeb. Bd. 1; Berlin: Springer-Verlag, 1932.
- [Rob] R.A.Robertello, *An invariant of knot cobordism*, Comm. Pure Appl. Math. 18 (1965), 543-555.
- [Rol] D.Rolfsen, *Knots and links*, Publish or Perish, Inc. Berkeley 1976, Math. Lect. Series 7.
- [Tsu] Added for e-print: T. Tsukamoto, The fourth skein module of 3-dimensional manifolds, PhD dissertation, George Washington University, May, 2000.
- [Wa] B.Wajnryb, *Markov classes in certain finite symplectic representations of braid groups*, these proceedings. Added for e-print: Braids (Santa Cruz, CA, 1986), 687-695, Contemp. Math., 78, Amer. Math. Soc., Providence, RI, 1988.
- [We] H.Wendt, *Die gordische Auflösung von Knoten*, Math. Z. 42 (1937), 680-696.

Department of Mathematics
 Warsaw University
 00901 Warszawa, PKiN IXp.
 Poland